

Lab: Millikan's Oil-Drop Experiment

AP Physics

Background

From 1909 to 1913, Robert Millikan performed a series of experiments designed to measure the charge of an electron. His general strategy was to place charges (electrons) on very small drops of oil, and then place those oil-drops in an electric field. By considering the Force of gravity, the electric Force, and the drag Force (air friction) acting on the drops, Millikan collected enough data that he was able to determine that the fundamental quantity of charge (the electron) is $-1.60 \times 10^{-19} \text{ C}$. For his experiments, Millikan won the Nobel Prize in Physics in 1923.

Learn more about Robert Millikan at <https://www.youtube.com/watch?v=sUc13Q8CF3s>, from *The Mechanical Universe* (YouTube).

Objectives

To observe the motion of charged particles in an electric field (a modified version of Millikan's experiment), and to perform a data analysis of information "collected" in a statistical simulation of Millikan's experiment.

Equipment

Millikan oil-drop device (set up in class)

Group "data" collected in a classroom simulation


Spreadsheet software (LibreOffice *Calc*, Apple *Numbers*, Google *Sheets*, or Microsoft *Excel*)

Procedure

Part A. Millikan's Device

1. Examine the device that has been set up for this lab.

A schematic diagram of Millikan's apparatus is shown here (from wikipedia.org):



2. Read this explanation of the device's basic operation

To determine the fundamental unit of electric charge (e^-), Millikan sprayed droplets of oil in to a chamber exposed to an electric field E . The droplets, which are charged by friction in the spraying process, experience an electrostatic force F_e according to the equation $F = qE$.

The drops also experience a downward force due to gravity, F_g , and a force of air friction F_{drag} when in motion. By analyzing the motion of the drops in the absence of the E field, it is possible to determine a droplet's mass m .

Lab: Millikan's Oil-Drop Experiment

AP Physics

With this same oil drop, if the Electric field is adjusted so that the droplet is suspended motionless, then the electrostatic force F_e will be just equal to the F_g .

$$F_e = F_g$$
$$qE = mg$$
$$q = \frac{mg}{E}$$

For a given oil-drop, there will be only one electric field that will suspend it. The strength of the field that supports a given oil drop depends on the mass of the droplet as well as the net charge of that droplet, and a given mass can have a wide variety of different charges, depending on how many electrons have been added to it (or subtracted from it) in the charging process.

How, then, did Millikan determine the magnitude of the fundamental electric charge? (Rhetorical question—we'll see how in the lab.)

3. Examine the online simulation of the oil drop experiment.

Click “Start” on the simulation at <https://www.magnus-karlsson.nu/millikan/> and see how gravity affects the oil drops as they appear in the Electric field. You can vary the strength of the Electric field by using the slider at the bottom of the simulation. Are you easily able to create a field that balances the force of gravity acting on that drop?

4. Draw a free-body diagram of the forces acting on an oil drop in static equilibrium in an Electric Field. Include field lines in your free-body diagram.

5. How did Millikan know what the Electric Field strength was? How could he vary it? In the graphic above and in the simulation you performed, you can see that the electric field is maintained between two conducting plates (one positive, one negative) with a potential difference between them. If a 500.0 Volt potential difference exists between the two plates, and there is a 1.50 centimeter distance between them, calculate the strength of the electric field between the plates.

Part B. Data Analysis

- 1. Get a data set of Electric field strengths and charge mass.**
Download the Common Separated Values (CSV) data at
<https://www.crashwhite.com/apphysics/materials/assignments/lab/millikan/data.csv>
- 2. Analyze a subset of that data to determine the fundamental quantity of charge, i.e. the charge of an electron.**
 - a. How will you calculate the electric field that each droplet has been exposed to, given the potential and gap distance specified in the data?
 - b. How will you calculate the charge on a given droplet, floating motionless in the electric field?
 - c. Use a spreadsheet to record a subset of the data values you downloaded, and use the spreadsheet's cell-calculation capabilities to identify the charge on each droplet.
 - d. Create a bar graph showing the charges of the different droplets. Are there any patterns immediately evident?
 - e. To help identify any patterns in your data, copy the column of charges that you've calculated and paste just the values (not the formulas) into a new column, then sort them. What does a bar graph of the charges look like now? Are there any patterns evident?
 - f. Discuss with the instructor additional ways to pull meaningful data from your results.